



RIVER RANCH
DENTAL

Post-Operative Instructions for Root Canal Therapy

Please follow the instructions outlined below, and contact us should any questions or problems arise.

1. DO NOT eat or drink anything for the next half hour. DO NOT try to feel around your tooth with your tongue. You have a temporary filling in the tooth and it takes about half an hour to harden.
2. **You need to have a permanent filling or a crown placed on your tooth within 1 MONTH of the root canal being completed.** It is essential for you to follow up with this treatment. If this is not done, the tooth is very likely to fracture or to develop new decay underneath the temporary filling which may cause your root canal to fail. *You will be responsible for all costs incurred if you fail to follow this instruction.*
3. DO NOT use the tooth to bite down on anything hard (peanuts, pretzels, ice, etc.) until the permanent filling/crown has been placed on the tooth. Again, the tooth is prone to fracture and if you bite down on anything too hard or crunchy you may crack the tooth.
4. It is normal for the temporary filling to “divot” in with use. It is very rare for it to fall out entirely. If the temporary falls out you should contact us, or your general dentist as soon as possible. If it happens when our office is closed, we recommend you purchase some temporary filling material from any pharmacy and place a dab in the tooth until you have an opportunity to see us at our office.
5. *Some minor discomfort in the area is normal following the root canal. It is normal for the tooth to be uncomfortable for 2-3 days after today’s treatment.* Sometimes, depending on the circumstances, the tooth and surrounding tissues may remain sore for **a few weeks** post treatment. The three most common reasons for pain are:
 - Sore jaw joint from having your mouth open for a prolonged time.
 - Sore muscle from the injection site.
 - Sore gum from the rubber dam placement.
6. WHAT DO I DO ABOUT THE PAIN??? All of the above scenarios should be handled with over the counter medication, primarily. We recommend you take:
600 mg Ibuprofen (3 over the counter pills of Advil or Motrin or generic equivalent) every 6 hours for the next 2 days.
OR
(if unable to take ibuprofen)
1000mg Acetaminophen (2 extra strength Tylenol or generic equivalent) every 6-8 hours for the next 2 days.
90% of the time this is enough to handle the discomfort. If it is not, please contact our office and we will ensure you are comfortable.
7. If you are given any prescription medications related to this treatment please take them as instructed by your doctor.
8. You may floss and brush your tooth as normal, unless told otherwise by the doctor

